




# HPV

## (16/18 genotypes) DNA Test (Real-Time PCR)

**HPV (16/18 genotypes) DNA Test (Real-Time PCR)** is used to qualitatively detect human papillomavirus (HPV) type 16 and type 18 deoxyribonucleic acid (DNA) in cervical exfoliated cells and genitourinary secretions specimens.


# FEATURES



# BACKGROUND

- Cervical cancer is the fourth most common cancer among women globally, with an estimated **604,000** new cases and **342,000** deaths in 2020.
- More than 95% of cervical cancer is due to the human papillomavirus (HPV). HPV may repeatedly infect people and gradually become chronic and pre-cancerous lesions.
- Two HPV types (16 and 18) cause at least 70% of cervical cancers and are responsible for 50% of high-grade cervical pre-cancers.
- Cervical cancer can be cured if diagnosed at an early stage and treated promptly. 70% of women should be screened with a high-performance test by 35 years of age according to the new Global strategy adopted by World Health Assembly.

Estimated age-standardized incidence rates (World) in 2020, cervix uteri, all ages



All rights reserved. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization / International Agency for Research on Cancer concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate borderlines for which there may not yet be full agreement.

Data source: GLOBOCAN 2020  
Graph production: IARC  
(<http://gco.iarc.fr/today>)  
World Health Organization



# SCREENING TEST OF CERVICAL CANCER




To reach the 2030 targets of cervical cancer elimination, WHO now encourages countries to use HPV tests for cervical screening, including HPV DNA and HPV mRNA tests.

- HPV-DNA testing detects high-risk strains of HPV, which cause almost all cervical cancers.
- HPV mRNA detects HPV infections leading to cellular transformation.



Reference: <https://www.who.int/news-room/fact-sheets/detail/cervical-cancer>  
<https://www.who.int/news/item/12-04-2022-meeting-of-who-director-general-s-expert-group-on-cervical-cancer-elimination-19-april-2022>

## HPV PCR TESTING WORKFLOW



## PARAMETER

<b>Test principle</b>	Qualitative real-time fluorescent PCR
<b>Specimen type</b>	Cervical exfoliated cells, genitourinary secretions
<b>Specimen extraction method</b>	Spin column (recommended)
<b>Fluorescence channel</b>	FAM (HPV16/18 gene detection) VIC/HEX (Internal reference gene detection)
<b>LOD</b>	1.0E+04 copies/reaction
<b>Storage and shelf life</b>	Nucleic acid extraction reagent: 18-25°C, 9 months Nucleic acid amplification reagent: -20±5°C, 9 months Proteinase K and Carrier RNA: -20±5°C, 9 months
<b>Applicable Instruments</b>	ABI Prism 7500, ABI Prism 7300, LightCycler 480
<b>Certificate</b>	NMPA, FSC, CE

## MAIN COMPONENTS



## ORDER INFORMATION

Description	Cat. No.	Package
HPV(16/18 genotypes) DNA Test (Real-Time PCR)	DA0540	48 tests/kit

## CONTACT US

### Daan Gene Co., Ltd.

#### CHINA (Headquarters)

Add.: No.19, Xiangshan Road, Science Park, High & New Technology  
 Development District, Guangzhou, Guangdong, P.R. China  
 Tel.: +86-20-32068126  
 Fax: +86-20-32068352  
 Email: marketing@daangene.com

#### CANADA office

Add.: #413-4538 Kingsway, Burnaby, BC V5H 4T9, Canada  
 Tel.: +1604 451 7588  
 Fax: +604451.7587  
 Email: gbai@daangene.net

#### COLOMBIA office

Add.: Cl 93 No.16 46 of 702, Bogotá D.C.  
 Tel.: +57 3103940417  
 Email: alonso.zheng@daangene.net

Website: <https://en.daangene.com/>

